


Ecoregions of North Carolina




- | | |
|--|---|
| 45 Piedmont | 65 Southeastern Plains |
| 45a Southern Inner Piedmont | 65c Sand Hills |
| 45b Southern Outer Piedmont | 65l Atlantic Southern Loam Plains |
| 45c Carolina Slate Belt | 65m Rolling Coastal Plain |
| 45e Northern Inner Piedmont | 65p Southeastern Floodplains and Low Terraces |
| 45f Northern Outer Piedmont | |
| 45g Triassic Basins | 66 Blue Ridge |
| 45i Kings Mountain | 66c New River Plateau |
| 63 Middle Atlantic Coastal Plain | 66d Southern Crystalline Ridges and Mountains |
| 63b Chesapeake-Pamlico Lowlands and Tidal Marshes | 66e Southern Sedimentary Ridges |
| 63c Nonriverine Swamps and Peatlands | 66g Southern Metasedimentary Mountains |
| 63d Virginian Barrier Islands and Coastal Marshes | 66i High Mountains |
| 63e Mid-Atlantic Flatwoods | 66j Broad Basins |
| 63g Carolinian Barrier Islands and Coastal Marshes | 66k Amphibolite Mountains |
| 63h Carolina Flatwoods | 66l Eastern Blue Ridge Foothills |
| 63n Mid-Atlantic Floodplains and Low Terraces | 66m Sauratown Mountains |

PRINCIPAL AUTHORS: Glenn E. Griffith (NRCS), James M. Omernik (USEPA), Jeffrey A. Comstock (Indus Corporation), Michael P. Schafale (NCDENR), W. Henry McNab (USFS), David R. Lenat (NCDENR), and Trish F. MacPherson (NCDENR).

COLLABORATORS AND CONTRIBUTORS: James E. Harrison (USEPA), David L. Penrose (NCDENR), Roy L. Vick, Jr. (NRCS), Gerard McMahon (USGS), Robert Peet (UNC), Chip Smith (NRCS) and Tom Loveland (USGS).

CITING THIS MAP: Griffith, G.E., Omernik, J.M., Comstock, J.A., Schafale, M.P., McNab, W.H., Lenat, D.R., and MacPherson, T.F., 2002, Ecoregions of North Carolina, U.S. Environmental Protection Agency, Corvallis, OR, (map scale 1:1,500,000).

— Level III ecoregion County boundary —
 - - - Level IV ecoregion State boundary - - -



Ecoregions denote areas of general similarity in ecosystems and in the type, quality, and quantity of environmental resources. They are designed to serve as a spatial framework for the research, assessment, management, and monitoring of ecosystems and ecosystem components. Ecoregions are directly applicable to many state agency activities, including the selection of regional stream reference sites, the development of biological criteria and water quality standards, and the establishment of management goals for nonpoint-source pollution. They are also relevant to integrated ecosystem management, an ultimate goal of many federal and state resource management agencies.

The approach used to compile this map of North Carolina is based on the premise that ecological regions are hierarchical and can be identified through the analysis of the spatial patterns and the composition of biotic and abiotic phenomena that affect or reflect differences in ecosystem quality and integrity (Wiken 1986; Omernik 1987, 1995). These phenomena include geology, physiography, vegetation, climate, soils, land use, wildlife, and hydrology. The relative importance of each characteristic varies from one ecological region to another regardless of the hierarchical level. A Roman numeral hierarchical scheme has been adopted for different levels of ecological regions. Level I and Level II divide the North American continent into 15 and 52 regions, respectively (Commission for Environmental Cooperation Working Group 1997). At Level III, the continental United States contains 104 regions (United States Environmental Protection Agency [U.S. EPA] 2000). Level IV is a further subdivision of the Level III ecoregions. Explanations of the methods used to define the U.S. EPA's ecoregions are given in Omernik (1995), Omernik and others (2000), Griffith and others (1994, 1997), and Gallant and others (1989).

The Level III and IV ecoregions were compiled at a scale of 1:250,000 and depict revisions and subdivisions of earlier level III ecoregions that were originally compiled at a smaller scale (U.S. EPA 2000; Omernik 1987). Compilation of this map is part of a collaborative project primarily between the U.S. Department of Agriculture's Natural Resources Conservation Service (NRCS), the U.S. EPA National Health and Environmental Effects Research Laboratory (NHEERL), U.S. EPA Region IV, and the North Carolina Department of Environment and Natural Resources. This project is also associated with an interagency effort to develop a common framework of ecological regions (McMahon and others 2001). Regional collaborative projects, such as this one in North Carolina where some agreement can be reached among multiple resource management agencies, are a step in the direction of attaining commonality and consistency in ecoregion frameworks for the entire nation.

Comments regarding the Level III and IV Ecoregions of North Carolina map should be addressed to Glenn Griffith, USDA-NRCS, 200 SW 35th Street, Corvallis, OR 97333, (541) 754-4465, FAX: (541) 754-4716, email: griffith.glenn@epa.gov, or to James Omernik, USGS, 200 SW 35th Street, Corvallis, OR 97333, (541) 754-4458, email: omernik.james@epa.gov.

Literature Cited:

Commission for Environmental Cooperation Working Group, 1997, Ecological regions of North America - toward a common perspective: Montreal, Quebec, Commission for Environmental Cooperation, 71 p.

Gallant, A.L., Whittier, T.R., Larsen, D.P., Omernik, J.M., and Hughes, R.M., 1989, Regionalization as a tool for managing environmental resources: Corvallis, Oregon, U.S. Environmental Protection Agency EPA/600/3-89/060, 152 p.

Griffith, G.E., Omernik, J.M., Wilton, T.F., and Pierson, S.M., 1994, Ecoregions and subregions of Iowa - a framework for water quality assessment and management: The Journal of the Iowa Academy of Science, v. 101, no. 1, p. 5-13.

Griffith, G.E., Omernik, J.M., and Azevedo, S.H., 1997, Ecoregions of Tennessee: Corvallis, Oregon, U.S. Environmental Protection Agency, National Health and Environmental Effects Research Laboratory, EPA/600/R-97/022, 51 p.

McMahon, G., Gregonis, S.M., Waltman, S.W., Omernik, J.M., Thorson, T.D., Freeouf, J.A., Rorick, A.H., and Keys, J.E., 2001, Developing a spatial framework of common ecological regions for the conterminous United States: Environmental Management, v. 28, no. 3, p. 293-316.

Omernik, J.M., 1987, Ecoregions of the conterminous United States (map supplement): Annals of the Association of American Geographers, v. 77, no. 1, p. 118-125, scale 1:7,500,000.

Omernik, J.M., 1995, Ecoregions-a spatial framework for environmental management, in Davis, W.S. and Simon, T.P., eds., Biological assessment and criteria-tools for water resource planning and decision making: Boca Raton, Florida, Lewis Publishers, p. 49-62.

Omernik, J.M., Chapman, S.S., Lillie, R.A., and Dumke, R.T., 2000, Ecoregions of Wisconsin: Transactions of the Wisconsin Academy of Sciences, Arts and Letters, v. 88, no. 2000, p. 77-103.

U.S. Environmental Protection Agency, 2000, Level III ecoregions of the continental United States (revision of Omernik, 1987): Corvallis, Oregon, U.S. Environmental Protection Agency-National Health and Environmental Effects Research Laboratory, Map M-1.

Wiken, E., 1986, Terrestrial eozones of Canada: Ottawa, Environment Canada, Ecological Land Classification Series no. 19, 26 p.